



PRESS RELEASE

ASN Contacts:

Christine Feheley (202) 640-4638 | cfeheley@asn-online.org

Tracy Hampton thampton@nasw.org

STUDY EXAMINES QUALITY OF LIFE IN PATIENTS WITH KIDNEY DISEASE IN INDIA

Even early stages of the disease can have negative impacts on mental and physical health.

Highlights

- Between 15 and 22 out of every 100 patients in India with mild-to-moderate chronic kidney disease had significant impairment in at least 1 of the 5 domains of quality of life.
- Quality of life scores were associated with sociodemographic factors (lower income, poor education, and female gender), with almost no major impact of medical- or disease-related variables.

Washington, DC (January 30, 2020) — A new study indicates that even early stages of chronic kidney disease (CKD) can negatively impact individuals' quality of life. The findings, which appear in an upcoming issue of *CJASN*, point to the importance of addressing, in addition to the medical aspects of chronic diseases, other factors that are important to patients.

A common perception of CKD is that patients are generally well until they require dialysis or a kidney transplant, but even patients with early stages of the disease may struggle with medical issues and declining quality of life. To provide a more accurate assessment of the quality of life in adults in India with mild-to-moderate CKD, a team led by Vivekanand Jha, MD (George Institute for Global Health) and Gopesh Modi, MD (Samarpan Kidney Institute and Research Center) examined data from the Indian Chronic Kidney Disease Study.

“There is almost no information on quality of life measurements in patients with CKD from low-middle income countries including India, which is home to about one-sixth of the world population and has over a million people with CKD,” said Dr. Jha. “The Indian Chronic Kidney Disease Study is the only large, multicentric prospective cohort of patients with CKD representing all parts of India. The primary objective of the study is to examine the natural history of CKD and the association of sociodemographic, clinical, laboratory, and biological variables with clinical outcomes, including those important to patients.”

In the study, all patients had their quality of life assessed at baseline and annually. This included assessments of mental health, physical health, burden of kidney disease, symptoms and problems of kidney disease, and effects of kidney disease. “We found that between 15 and 22 out of every 100 Indian patients with mild-to-moderate CKD had significant impairment in at least 1 of the 5 domains of quality of life,” Dr. Jha said.

After adjusting for confounding factors, lower quality of life scores were associated with lower income, poor education, and female gender. “The findings point to the need to prioritize interventions targeted to these populations. Better understanding of the nature of these associations, and how they evolve over time—which will be ascertained in the follow-up phase of the study—will allow development of tailored interventions.”

An accompanying editorial assesses the study's findings.

Study co-authors include Ashok Yadav, PhD, Arpita Ghosh, PhD, Kajal Kamboj, MSc, Prabhjot Kaur, PhD, Vivek Kumar, MD, Shobhit Bhansali, PhD, Narayan Prasad, MD, Manisha Sahay, MD, Sreejith Parameswaran, MD, Santosh Varughese, MD, Sishir Gang, MD, Shivendra Singh, MD, Dipankar Sircar, MD, N Gopalakrishnan, MD, Ajay Jaryal, MD, Samjay Vikrant, MD, and Seema Baid-Aggarwal, MD.

Disclosures: The authors reported no financial disclosures.

The article, entitled “Non-Medical Factors and Health-Related Quality of Life in CKD in India,” will appear online at <http://cjasn.asnjournals.org/> on January 30, 2020, doi: 10.2215/CJN.06510619.

The editorial, entitled “Socioeconomic Determinants of Quality of Life in Patients with Kidney Diseases,” will appear online at <http://cjasn.asnjournals.org/> on January 30, 2020, doi: 10.2215/CJN.14941219.

The content of this article does not reflect the views or opinions of The American Society of Nephrology (ASN). Responsibility for the information and views expressed therein lies entirely with the author(s). ASN does not offer medical advice. All content in ASN publications is for informational purposes only, and is not intended to cover all possible uses, directions, precautions, drug interactions, or adverse effects. This content should not be used during a medical emergency or for the diagnosis or treatment of any medical condition. Please consult your doctor or other qualified health care provider if you have any questions about a medical condition, or before taking any drug, changing your diet or commencing or discontinuing any course of treatment. Do not ignore or delay obtaining professional medical advice because of information accessed through ASN. Call 911 or your doctor for all medical emergencies.

Since 1966, the American Society of Nephrology (ASN) has been leading the fight to prevent, treat, and cure kidney diseases throughout the world by educating health professionals and scientists, advancing research and innovation, communicating new knowledge, and advocating for

the highest quality care for patients. ASN has more than 21,000 members representing 131 countries. For more information, visit www.asn-online.org.

###

Twitter: Study examines quality of life in patients with kidney disease in India.
George Institute (@GeorgeInstIN); Vivekanand Jha (@vjha126)

Facebook: A new study indicates that even early stages of chronic kidney disease (CKD) can negatively impact individuals' quality of life. The findings, which appear in *CJASN*, point to the importance of addressing more than the medical aspects of chronic diseases.

Media: Kannan Krishnaswamy – kkrishnaswamy@georgeinstitute.org.in